

## Karyological Evidence for a Cryptic Species of Piranha within *Serrasalmus rhombeus* (Characidae, Serrasalminae) in the Amazon

C. NAKAYAMA, M. JÉGU, J.I.R. PORTO, AND E. FELDBERG

Cytogenetic data obtained for *Serrasalmus rhombeus* revealed two cytotypes (cyto-type 1 and cyto-type 2) at four sampling sites. Cyto-type 1 consists of  $2n = 60, 20M + 24SM + 6ST + 10A$ , and was detected in specimens from Negro River (Anavil-hanas Islands), Solimões River (Camaleão Lake), confluence of Negro and Solimões rivers (Catalão Lake), and Uatumã River. Cyto-type 2 consists of  $2n = 58, 22M + 24SM + 2ST + 10A$  and was detected in specimens from Catalão and Camaleão Lakes. In Cyto-type 2, the first chromosome pair was almost twice the size of second pair, and the constitutive heterochromatin pattern of each cyto-type is distinct. The two cytotypes occurred sympatrically at Catalão and Camaleão lakes and no inter-mediate between the cytotypes were detected, suggesting each cyto-type represents a different fish species.

**M**ANY examples of cytogenetic variation have been reported in Neotropical fishes. For example, intraspecific cytogenetic variation in the diploid number, karyotype structures or C-banding patterns occur in nominal forms of *Hoplias malabaricus*, *Astyanax scabripinnis*, *Eigenmannia virescens*, and *Callichthys callichthys*. De-bates remain concerning the status of the cyto-types within these nominal species (Oliveira et al., 1988; Porto et al., 1992; Almeida-Toledo, 1998). In the Amazon, intraspecific cytotypic variation has been reported in *Callichthys callichthys* (Porto and Feldberg, 1993), *Plagioscion* sp. (Feldberg et al., 1999) and *Serrasalmus spilopleura* (Centofante, 2000; Nakayama et al., 2000). However, only a few studies have associated both morphological and karyotypic difference (Moreira Filho and Bertollo, 1991).

The genus *Serrasalmus* is distributed through-out the tropical regions of South America east-ward from the Andean Cordillera. The taxonomy and systematics of several species of this genus are not well known because most analyses were based on small sample sizes usually consisting of immature specimens (Machado-Allison and Fink, 1996). Thus, the number of recognized species continues to fluctuate (Merckx et al., 2000). *Serrasalmus rhombeus* is considered to be one of the most common piranhas in the Amazon. It was described from Surinam but is widespread throughout the Amazon Basin, Orinoco and Guiana coastal basins. Its pigmentation changes with age, being silvery with small dark spots when young (known as white piranha) and becoming progressively darker when adult (when they are known as black piranha or red-eyed piranha). According to Géry

(1976) *S. rhombeus* is a species complex (*rhombeus* group) comprising six to nine species (*S. rhombeus*, *S. albus*, *S. spilopleura*, *S. marginatus*, *S. brandtii*, *S. sanchezi* plus probably *S. gibbus*, *S. humeralis* and *S. pingke* = *S. elongatus*) that are recognized by a deep and/or compressed body. The aim of this paper is to document the cyto-genetic variation found in nominal forms of *S. rhombeus* collected in Central Amazon near Manaus, Amazonas, Brazil and to focus on a particular taxonomic problem that may be assisted by chromosomal analyses.

### MATERIALS AND METHODS

A total of 71 specimens of the nominal *S. rhombeus* were collected in the Central Amazon Basin between 25 March 1984 and 30 September 1996 at the following sites: (1) 45 specimens (17 males and 28 females) at the confluence of Negro and Solimões rivers (Catalão Lake); (2) 16 specimens (nine males and seven females) on Solimões River (Camaleão Lake-Marchantaria Island); (3) two specimens (two females) in the Uatumã River; (4) six specimens (two males and four females) in the Jatapu River; (5) one female specimen at Capu-capu River; (6) one female specimen at Anavil-hanas in the Negro River (Fig. 1).

Mitotic chromosomes, nucleolar organizing regions (NORs), and C-band were obtained through the methods of Bertollo et al. (1978), Howell and Black (1980), and Sumner (1972), respectively. The chromosomes were analyzed by measuring the short arm length, long arm length, and total length, by means of a dry-tip compass and a calliper. Mean values were calculated for each chromosome pair. The relative length (RL%) of each chromosome pair in relation to the total length of the haploid chromosome



TABLE 1. THE MAIN CYTOGENETIC DATA OF *Serrasalmus rhombeus*. (NORs = nucleolar organizer regions, n = number of specimens, NCell = number of cells analysed, 2n = diploid number, FN = Fundamental number, M = metacentric, SM = submetacentric, ST = subtelocentric, pSTA = short arm of chromosomes subtelocentric, Loc = Location).

Collecting sites	n	NCell	2n	NF	Formula			NORs		
					M	SM	ST	A	Range	Loc
CYTOTYPE 1										
Catalão Lake	10	519	60	110	20	24	06	10	5-12	pST-A
Camaleão Lake	09	313	60	110	20	24	06	10	5-12	pST-A
Uatumã River	02	64	60	110	20	24	06	10	5-12	pST-A
Jatapu River	06	268	60	110	20	24	06	10	5-12	pST-A
Capucapu River	01	26	60	110	20	24	06	10	5-12	pST-A
Anavilhanas Archip.	01	44	60	110	20	24	06	10	5-12	pST-A
CYTOTYPE 2										
Catalão Lake	35	1439	58	106	30	16	02	10	5-12	pST-A
Camaleão Lake	07	322	58	106	30	16	02	10	5-12	pST-A

larities observed among cytotypes encompass the proximal blocks on the long arms of one metacentric pair (seventh in both cytotype 1 and cytotype 2), the short arms totally hetero-chromatic of one submetacentric pair (14th in the cytotype 1 and 15th in the cytotype 2) and the centromeric blocks of acrocentric chromosomes. Because NORs were detected in acrocentric chromosomes, the NORs sites were considered to be positive for C-banding.

All specimens are morphologically very similar, however, specimens with cytotype 2 are smaller than those with cytotype 1. Individuals of cytotype 2 never exceeded 150 mm in standard length (SL), and females had ripe ovaries with 100 mm SL. Individuals with cytotype 1 exceeded 250 mm in SL and females have ripe ovaries with 160 mm SL.

#### DISCUSSION

This is the first report of karyotypes on specimens tentatively identified as *S. rhombeus*, and the chromosomal data presented here are part of a multidisciplinary study that encompasses parasitological, morphological, and molecular analyses on piranhas from the Amazon. In the Amazon floodplain near Manaus, *S. rhombeus* has two cytotypes that differ in diploid numbers, karyotypic formulae, and in C-banding patterns. It is noteworthy that cytotype 1 is the most widespread form being found in Uatumã, Negro and Solimões Rivers drainages, whereas cytotype 2 occurs only in the lower Negro River at the confluence with Solimões River. Because there is no cytological evidence of hybridization between the cytotypes (no intermediate

cytotypes) where they co-exist (Catalão and Camaleão lakes) the cytotypes appear to represent distinct species.

Besides chromosomal character each cytotype (species) is uniquely defined by a set of morphological and parasitological character states. Preliminary morphometric and meristic data (including number of supraneural bones) indicate subtle differences between the two species (A. Merckx and M. Jégu, unpubl.). We also observed that adults from cytotype 1 are considerably larger than those from cytotype 2. Monogenea parasite data also support to the hypothesis that they might be distinct species because individuals representing the two cytotypes have different gill parasite species within the monogenoid genera *Anachantorus* (Kritsky et al., 1992), *Mymarothecium* and *Notothecium* (Kritsky et al., 1996), *Amphitecium* (Kritsky et al., 1997), *Notothecium* and *Enalothecium* (Kritsky et al., 1998).

Examples of sympatric piranha species that possess a deep and/or compressed body typical of *S. rhombeus*-species complex seem not to be a rare event in Amazon river drainages: (1) *S. rhombeus* and *S. geryi* from Tocantins River (Jégu and Santos, 1988); (2) *S. altispinis* and *S. rhombeus* from Uatumã River (Merckx et al., 2000); and (3) cytotype 1 and cytotype 2 from lower Negro River (this work). Particularly in the latter example, external morphological characters do not help to differentiate the species and a reasonable explanation might be a recent speciation. Additional studies are necessary to understand why individuals with cytotype 2 present such a limited distribution as well as to test the hypothesis whether all species included in the *S. rhombeus*-species complex are a monophyletic unit.

## MATERIAL EXAMINED

22 of *S. rhombeus* (2n=58) were selected and deposited at the ichthyology collection of the Instituto Nacional de Pesquisas da Amazônia, Manaus, Amazonas, Brazil with the following accession numbers: INPA 7352-7356 (2n = 60) and INPA 7345-7351 (2n = 58).

## ACKNOWLEDGEMENTS

We thank L.A.C. Bertollo (Universidade Federal de São Carlos), V. Thatcher, and A. S. Teixeira (Instituto Nacional de Pesquisas da Amazônia), and three anonymous reviewers for their contribution on preliminary versions of this paper. This work was supported by the Brazilian (CNPq and INPA) and the French (IRD) research institutions.

## LITERATURE CITED

- ALMEIDA-TOLEDO, L. F. 1998. Cytogenetic markers in neotropical freshwater fishes, P.583-588. *In*: Phylogeny and classification of neotropical fishes. L. R. Malabarba, R. R. Reis, R. P. Vari, Z. M. S. Lucena, and C. A. S. Lucena (eds.). EDIPUCRS, Porto Alegre, RS.
- BERTOLLO, L. A. C., C. S. TAKAHASHI, AND O. MOREIRA-FILHO. 1978. Cytotaxonomic considerations on *Hoplias lacerdae* (Pisces, Erythrinidae). *Braz. J. Genet.* 1:103-120.
- CENTOFANTE, L. 2000. Análise cariotípica no "complexo" *Serrasalmus spilopleura* (Characiformes, Serrasalminidae) da bacia amazônica central. Unpubl. M.Sc. Diss., Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus-AM.
- FELDBERG, E., J. I. R. PORTO, E. B. P. SANTOS, AND F. C. S. VALENTIM. 1999. Cytogenetic studies of two freshwater sciaenids of the genus *Plagioscion* (Perciformes, Sciaenidae) from the Central Amazon. *Genet. Mol. Biol.* 22:351-356.
- GÉRY, J. 1976. Les Genres de Serrasalminidae (Pisces-Characoidei). *Bull. Zool. Mus. Univ. van Amsterdam* 5:47-54.
- HOWELL, W. M., AND D. A. BLACK. 1980. Controlled silver staining of nucleolus organizer regions with a protective colloidal developer: a 1-step method. *Experientia* 36:1014-1015.
- JÉGU, M., AND G. M. SANTOS. 1988. Le genre *Serrasalmus* (Pisces, Characiformes) dans le Bas Tocantins (Brésil, Pará), avec la description d'une espèce nouvelle, *S. geryi*, du bassin Araguaia-Tocantins. *Rev. Hydrobiol. Trop.* 21:239-274.
- KRITSKY, D. C., W. A. BOEGER, AND L. R. EVERY. 1992. Neotropical Monogenoidea. 17. *Anacanthorus* Mizelle & Price, 1965 (Dactylogyridae, Anacanthorinae) from characid fishes of the central Amazon. *J. Helminthol. Soc. Wash.* 59:5-51.
- \_\_\_\_\_, \_\_\_\_\_, AND M. JÉGU. 1996. Neotropical Monogenoidea. 28. Ancyrocephalinae (Dactylogyridae) of piranha and their relatives (Teleostei, Serrasalminidae) from Brazil and French Guiana: species of *Notozothecium* Boeger and Kritsky, 1988, and *Mymarothecium* gen.n. *Ibid.* 63:153-173.
- \_\_\_\_\_, \_\_\_\_\_, AND \_\_\_\_\_. 1997. Neotropical Monogenoidea. 29. Ancyrocephalinae (Dactylogyridae) of piranha and their relatives (Teleostei, Serrasalminidae) from Brazil: species of *Amphithecium* Boeger and Kritsky, 1988, *Heterothecium* gen.n and *Pithanothecium* gen.n. *Ibid.* 64:25-54.
- \_\_\_\_\_, \_\_\_\_\_, AND \_\_\_\_\_. 1998. Neotropical Monogenoidea. 31. Ancyrocephalinae (Dactylogyridae) of piranha and their relatives (Teleostei, Serrasalminidae) from Brazil: species of *Notothecium* Boeger and Kritsky, 1988, and *Enallothecium* gen.n. *Ibid.* 65:31-49.
- LEVAN, A., K. FREDGA, AND A. A. SANDBERG 1964. Nomenclature for centromeric position on chromosomes. *Hereditas* 52:201-220.
- MACHADO-ALLISON, A., AND W. FINK 1996. Los peces caribes de Venezuela: diagnosis, claves, aspectos ecológicos y evolutivos. Universidad Central de Venezuela, Consejo de Desarrollo Científico y Humanístico, Caracas. Colección monografías 52.
- MERCKX, A., M. JÉGU, AND G. M. SANTOS. 2000. Une nouvelle espèce de *Serrasalmus* (Teleostei: Characidae: Serrasalminae), *S. altispinis* n. sp., décrite du rio Uatumã (Amazonas, Brésil) avec une description complémentaire de *S. rhombeus* (Linnaeus, 1766) du plateau Guyanais. *Cybum* 24:81-201.
- MOREIRA-FILHO, O., AND L. A. C. BERTOLLO. 1991. *Asytanax scabripinnis* (Pisces, Characidae): a species complex. *Brazil. J. Genet.* 14:331-357.
- NAKAYAMA, C. M., J. I. R. PORTO, AND E. FELDBERG. 2000. Ocorrência de dois citótipos em *Serrasalmus spilopleura* Kner, 1860 (Characiformes, Serrasalminidae) da região de confluência dos rios Negro e Solimões, Amazonas, Brasil. *Acta Amazonica* 30:149-154.
- OLIVEIRA, C., L. F. ALMEIDA-TOLEDO, F. FORESTI, H. A. BRITSKI, AND S. A. TOLEDO FILHO. 1988. Chromosome formulae of neotropical freshwater fishes. *Brazil. J. Genet.* 11:577-624.
- PORTO, J. I. R., E. FELDBERG, 1993. Is *Callichthys* Linné (Siluriformes, Callichthyidae) a monotypic genus? *Acta Amazonica* 24:311-314.
- \_\_\_\_\_, AND \_\_\_\_\_, C. M. NAKAYAMA, AND J. N. FALCÃO. 1992. A checklist of chromosome numbers and karyotypes of Amazonian freshwater fishes. *Rev. Hydrobiol. Trop.* 25:287-299.
- SUMNER, A.T. 1972. A simple technique for demonstrating centromeric heterochromatin. *Expl. Cell Res.* 75:304-306.
- (CMN, JIRP, EF) INPA, COORDENAÇÃO DE PESQUISAS EM BIOLOGIA AQUÁTICA, LABORATÓRIO DE GENÉTICA DE PEIXES, CX. POSTAL 478, 69011-970 MANAUS, BRASIL.; AND (MJ) ANTENNE IRD, LABORATOIRE D'ICHTHYOLOGIE, M.N.H.N., 43, RUE CUVIER F-75231 PARIS CEDEX 05 FRANCE. Email: (EF) feldberg@inpa.gov.br. Submitted: 22 March 2000. Accepted: 23 March 2001. Section editor: J.D. McEachran.